THE JOURNAL OF CHEMICAL THERMODYNAMICS

Volume 26, 1994

EXECUTIVE EDITOR

Professor M. L. McGlashan
Patoka, Fairwarp, Uckfield, East Sussex TN22 3DT, U.K.

EDITORS

Dr P. A. G. O'Hare
National Institute of
Standards and Technology
Building 222
Gaithersburg, MD 20899
U.S.A.

Professor W. A. Wakeham
Dept of Chem. Engineering
and Chem. Technology
Imperial College
London SW7 2BY
U.K.

Professor R. D. Weir
Dept of Chemistry and
Chem. Engineering
Royal Military College
Kingston
Ontario, Canada K7K 5L0

ADVISORY BOARD

Chairman: G. Somsen (The Netherlands)

G. Della Gatta (Italy)
J. H. Dymond (U.K.)
M. B. Ewing (U.K.)
J.-P. E. Grolier (France)
L. V. Gurvich (Russia)
Y. P. Handa (Canada)
H. F. Holmes (U.S.A.)
G. M. Olofsson (Sweden)
D. Patterson (Canada)
G. Pilcher (U.K.)
J. A. Rard (U.S.A.)
M. A. Ribeiro da Silva
(Portugal)
G. M. Schneider (Germany)
M. Sorai (Japan)
W. V. Steele (U.S.A.)
F. L. Swinton (U.K.)

ACADEMIC PRESS

London San Diego New York
Boston Sydney Tokyo Toronto

Copyright © 1994 Academic Press Limited

ALL RIGHTS RESERVED

No part of this volume may be reproduced in any form, by photostat, microfilm, or any other means, without written permission from the publishers.

Printed in Great Britain

CONTENTS OF VOLUME 26

Number 1, January 1994

Apelblat, A. Freezing-temperature depressions of aqueous potassium hydrogen carbonate 1

Guil, J. M.; Pérez Masiá, A.; Ruiz Paniego, A.; Trejo Menayo, J. M. Enthalpies of adsorption of hydrocarbons on discandium trioxide determined calorimetrically and from adsorption isotherms 5

Chyliński, K. Isothermal compressibilities and excess molar volumes of $\{x CH_3C(CH_3)_2CH_2CH(CH_3)CH_3 + (1-x)$*cis*-$C_{10}H_{18}\}$ at the temperature T = 298.15 K; and of $\{x CH_3C(CH_3)_2CH_2CH(CH_3)CH_3 + (1-x)$*trans*-$C_{10}H_{18}\}$ at T = 298.15 K and 318.15 K 15

Blanco, S. T.; Muñoz Embid, J.; Otín, S. Excess volumes of (2,2,4-trimethylpentane + methanol or ethanol or propan-2-ol or di-1-methylethyl ether or 1,1-dimethylethyl methyl ether) and of (methanol or ethanol or propan-2-ol + di-1-methylethyl ether or 1,1-dimethylethyl methyl ether) at the temperature 298.15 K 23

Castro, I.; Pintos, M.; Amigo, A.; Bravo, R.; Paz Andrade, M. I. Excess enthalpies of (tetrahydrofuran or tetrahydropyran + an *n*-alkane) at the temperature 298.15 K . 29

Zhu, S.; Shen, S.; Benson, G. C.; Lu, B. C.-Y. Excess enthalpies of (methyl 1,1-dimethylpropyl ether + an *n*-alkane) at the temperature 298.15 K 35

Gonzalez, E.; Ortega, J. Thermodynamic properties of (a methyl ester + an *n*-alkane) VI. H_m^E and V_m^E for $\{x CH_3(CH_2)_{u-1}CO_2CH_3$ (u = 1 to 6) $+ (1-x)CH_3(CH_2)_{14}CH_3\}$. 41

Apelblat, A. Enthalpies of solution of citrates and hydrogen citrates of lithium, sodium, and potassium 49

García, J.; López, E. R.; Fernández, J.; Amigo, A.; Paz Andrade, M. I. Intramolecular-proximity effect on the excess enthalpies of (a dichloroalkane + an alkan-2-one). 53

Korobov, M. V.; Sidorov, L. N. Thermodynamic properties of fullerenes 61

Letcher, T. M.; Domańska, U. The excess enthalpies of (acetonitrile + an ether) at the temperature 298.15 K 75

Acree, W. E., Jr.; Tucker, S. A.; Pilcher, G.; Toole, G. Enthalpies of combustion of 4-dimethylaminonitrosobenzene and 4-dimethylaminonitrobenzene 85

Kalinkin, A. M.; Vasin, S. K.; Antropova, O. V. Solubility diagram of (lithium sulfate + sodium sulfate + rubidium sulfate)(aq) at the temperature 298.15 K . 91

Wormald, C. J.; Lloyd, M. J. Excess molar enthalpies for $(0.5H_2O + 0.5CO_2)$ at temperatures from 598.2 K to 698.2 K and pressures up to 20 MPa 101

Advice to authors. 111

Number 2, February 1994

Letcher, T. M.; Domańska, U. Excess volumes of (acetonitrile + an ether) at the temperature 298.15 K 113

10 283HI XL 968
5/95 31249-200 NULE

Orge, B.; Iglesias, M.; Tojo, J.; Legido, J. L. Densities and refractive indices of $\{x_1CH_3COCH_3 + x_2CH_3OH + (1-x_1-x_2)C_6H_5Cl\}$ at the temperature 298.15 K . 121

Kabo, G. J.; Kozyro, A. A.; Marchand, A. P.; Diky, V. V.; Simirsky, V. V.; Ivashkevich, L. S.; Krasulin, A. P.; Sevruk, V. M.; Frenkel, M. L. Thermodynamic properties of heptacyclotetradecane $C_{14}H_{16}$ 129

Gospodinov, G. G. Heat capacities and derived thermodynamic properties of magnesium tellurates(IV) 143

Svoboda, V.; Čejka, D.; Zábranský, M. Heat capacities of six isomers of pentanol in the temperature range from 303 K to 326 K 147

Artigas, H.; Lafuente, C.; Rodríguez, V.; Royo, F. M.; Urieta, J. S. Excess volumes of (1-bromobutane + each of several isomers of butanol) at the temperatures 298.15 K and 313.15 K 151

Konings, R. J. M.; Hildenbrand, D. L. The vapor pressure, infrared spectrum, and thermodynamic properties of $ZrF_4(g)$ 155

Nagata, I.; Książczak, A. Excess enthalpies for (ethanol + propanone) and (ethanol + propanone + benzene) at the temperature 298.15 K 165

Wang, Z.-C.; Yu, H.-L.; Hu, Y.-F. Isopiestic studies on (mannitol + sorbitol + sucrose)(aq) at the temperature 298.15 K. Comparison with the partial-ideal-mixture model 171

Nikitin, E. D.; Pavlov, P. A.; Bessonova, N. V. Critical constants of *n*-alkanes with from 17 to 24 carbon atoms 177

Ferro, D.; Martino, R.; Della Gatta, G. Sublimation enthalpies and entropies of thiourea, 1-methyl-2-thiourea, 1,3-dimethyl-2-thiourea, and 1,1,3,3-tetramethyl-2-thiourea 183

Yu, C.-H.; Tsai, F.-N. Excess volumes of (tetralin + an *n*-alkan-1-ol) 191

Kumar, P.; Bhaisare, M. P.; Kudchadker, A. P. Excess molar volumes of $[xCH_3(CH_2)_4OH + (1-x)\{CH_3COCH(CH_3)CH_3$ or $CH_3COCH_2CH(CH_3)CH_3$ or $CH_3CH(OH)CH(CH_3)CH_3$ or $CH_3CH(CH_3)OCOCH_3$ or $CH_3CH_2CH(CH_3)$-$OCOCH_3$ or $CH_3(CH_2)_2CH(CH_3)OCOCH_3$ or $CH_3CH(CH_3)CH(CH_3)CH_3\}]$ at temperatures from 288.15 K to 328.15 K and each at saturation pressure . . 197

Zeng Wenming; Chen Qiyuan; Chen Xinmin Determination of the standard molar enthalpy of formation of the ion $Al(OH)_4^-(aq)$ 205

Kume, Y.; Muraoka, H.; Matsuo, T.; Suga, H. Low-temperature heat capacities of ammonium hexachloroselenate and of its deuterated analogue 211

BOOK REVIEW (**Authors**, Title, **Reviewer**)

Kubaschewski, O.; Alcock, C. B.; Spencer, P. J. Materials Thermochemistry. Sixth edition, Revised. **J. G. Edwards** 223

Number 3, March 1994

Wood, R. H.; Carter, R. W.; Quint, J. R.; Majer, V.; Thompson, P. T.; Boccio, J. R. Aqueous electrolytes at high temperatures: comparison of experiment with simulation and continuum models. (Being the 41st Huffman Memorial Award Lecture of the Calorimetry Conference delivered on 19 July 1993 at its 48th annual Conference held at Durham, NC, U.S.A.) 225

Cao, H.; Boerio-Goates, J. Heat-capacity measurements and thermodynamic functions of langbeinite-type $K_2Cd_2(SO_4)_3$ and $K_2Ca_2(SO_4)_3$ at temperatures from 10 K to 340 K 251

Gampe, T.; Libuś, Z. Osmotic and activity coefficients of $C_2H_5SO_4Na$(aq) and $C_2H_5SO_4K$(aq) at the temperature 298.15 K 261

Holmes, H. F.; Busey, R. H.; Simonson, J. M.; Mesmer, R. E. $CaCl_2$(aq) at elevated temperatures. Enthalpies of dilution, isopiestic molalities, and thermodynamic properties 271

Tomaszkiewicz, I.; Passmore, J.; Schatte, G.; Sutherland, G. W.; O'Hare, P. A. G. Thermochemistry of octasulfur-bis-hexafluoroarsenate $S_8(AsF_6)_2$ containing the lattice-stabilized S_8^{2+} 299

Smruti Dash; Ziley Singh; Prasad, R.; Sood, D. D. The standard molar Gibbs free energy of formation of $SrZrO_3$(cr) 307

Ribeiro da Silva, M. A. V.; Ferrão, M. L. C. C. H. Enthalpy of sublimation of bis(pentane-2,4-dionato)copper(II) 315

Ohta, T.; Yamamuro, O.; Suga, H. Heat capacity and enthalpy of sublimation of sulfur hexafluoride 319

Fonseca, I. M. A.; Lobo, L. Q. (Vapour + liquid) equilibria of (ammonia + xenon) at the temperature 195.48 K 333

Number 4, April 1994

Torres, L. A.; Pérez, A; Farfán, N.; Castillo D.; Santillan, R. L. Rotating-bomb combustion calorimetry and the standard enthalpies of formation of two borinic esters 337

Simonson, J. M.; Oakes, C. S.; Bodnar, R. J. Densities of NaCl(aq) to the temperature 523 K at pressures to 40 MPa measured with a new vibrating-tube densitometer . 345

Sahayam, M.; Hedwig, G. R. Enthalpies of dilution of aqueous solutions of the amino acids L-threonine and L-asparagine at the temperature 298.15 K 361

Fujimori, H.; Oguni, M. Calorimetric study of D,L-propene carbonate: observation of the β- as well as α-glass transition in the supercooled liquid. 367

Pawlak, Z.; Wisniewska, M.; Richert, M. Homoconjugation and acidity constants of protonated trimethyl-*N*-oxide and of picric acid in (methanol + acetonitrile) . 379

Gilgen, R.; Kleinrahm, R.; Wagner, W. Measurement and correlation of the (pressure, density, temperature) relation of argon. I. The homogeneous gas and liquid regions in the temperature range from 90 K to 340 K at pressures up to 12 MPa . . 383

Gilgen, R.; Kleinrahm, R.; Wagner, W. Measurement and correlation of the (pressure, density, temperature) relation of argon. II. Saturated-liquid and saturated-vapour densities and vapour pressures along the entire coexistence curve. 399

Zhu, S.; Shen, S.; Benson, G. C.; Lu, B. C.-Y. Excess enthalpies of (2,4-dimethyl-3-oxapentane + one or more *n*-alkanes) at the temperature 298.15 K. Part I: Binary mixtures 415

Zhu, S.; Shen, S.; Benson, G. C.; Lu, B. C.-Y. Excess enthalpies of (2,4-dimethyl-3-oxapentane + one or more *n*-alkanes) at the temperature 298.15 K. Part II: Ternary mixtures 421

Hu, Y.-F.; Wang, Z.-C. Isopiestic studies on {NaCl (m_B) + NH_4Cl (m_C) + $BaCl_2$ (m_D)}(aq) at the temperature 298.15 K. A quaternary system obeying Zdanovskii's rule 429

Settle, J. L; Greenberg, E.; Hubbard, W. N.; O'Hare, P. A. G. The standard molar enthalpy of formation of liquid hydrogen fluoride $\Delta_f H^\circ_m$(HF, l) at the temperature 298.15 K determined by direct combination of the elements in a fluorine-bomb calorimeter 435

Number 5, May 1994

Yang, X. N.; Zhang, J. H. Excess enthalpies of binary mixtures containing cyclohexanone 449

Wang Haijun; Zheng Guokang; Chen Mingzhi Excess volumes of (acrylonitrile + an aromatic hydrocarbon) at the temperature 298.15 K 457

An Xueqin; Shen Weiguo The (liquid + liquid) critical phenomena of (a polar liquid + an *n*-alkane). II. Coexistence curves of (*N*,*N*-dimethylacetamide + octane) . . 461

Chirico, R. D.; Gammon, B. E.; Hossenlopp, I. A.; Steele, W. V. Heat capacities of 1,2,3,4-tetrahydrophenanthrene at temperatures between 5 K and 430 K and revised heat capacities of 9,10-dihydrophenanthrene at temperatures between 10 K and 350 K 469

Pawlak, Z.; Kuna, S.; Richert, M.; Giersz, E.; Liwo, A.; Chmurzynski, L. Proton transfer and heteroconjugation of ammonium ions with *N*-bases in cyclohexanone, propanone, and butan-2-one 483

Aune, R. E.; Sridhar, S.; Du Sichen A galvanic-cell study of (nickel + tungsten + oxygen) in the temperature range 1034 K to 1317 K 493

Midura, J.; Pomianek, T. Activity of Pb in the alloy: $\{(1-x)Cu+xPb\}(l)$ . . . 507

Steele, W. V.; Chirico, R. D.; Nguyen, A.; Knipmeyer, S. E. The thermodynamic properties of 2-methylaniline and *trans*-(R,S)-decahydroquinoline . . . 515

Nagata, I. (Liquid + liquid) equilibria and excess enthalpies for (methanol + benzene or propanone + hexane) at the temperature 298.15 K 545

Letcher, T. M.; Domańska, U. The excess enthalpies of (tributylamine + an ether) at the temperature 298.15 K 553

Number 6, June 1994

Jett, M.; Fleyfel, F.; Kobayashi, R. The nature of (p,V_m,T) surfaces of mixtures: (quantum + normal) and (normal + hydrogen-bonded), as summarized by their isochoric inflection loci. 561

Letcher, T. M.; Bayles, J. W.; Moollan, W. C. The determination of activity coefficients at infinite dilution using g.l.c. with moderately volatile solvents (*cis*- and *trans*-decalin) at the temperature 298.15 K 571

Lavut, E. G.; Chelovskaya, N. V.; Belysheva, G. A.; Demin, V. N.; Zlomanov, V. P. Enthalpy of formation of indium telluride InTe 577

Holmes, H. F.; Mesmer, R. E. An isopiestic study of $\{(1-y)NaHSO_4 + yNa_2SO_4\}(aq)$ at elevated temperatures 581

Rodriguez, V.; Artigas, H.; Santafé, J.; Royo, F. M.; Urieta, J. S. Excess volumes of (an alkanol + a haloalkane) at the temperatures 298.15 K and 313.15 K . . . 595

Klein, R.; Jacob, I.; O'Hare, P. A. G.; Goldberg, R. N. Solution-calorimetric determination of the standard molar enthalpies of formation of the pseudobinary compounds $Zr(Al_xFe_{1-x})_2$ at the temperature 298.15 K 599

Artal, M.; Muñoz Embid, J.; Velasco, I. Excess enthalpies and excess volumes of (1-bromobutane or 1-iodobutane + 1-chloropropane or 1-chloropentane) and of (1-iodobutane + bromoethane or 1-bromobutane) at the temperature 298.15 K . 609

McElroy, P. J.; Ji Fang Compression factors and virial coefficients of (methane + ethane) 617

Hasch, B. M.; Maurer, E. J.; Ansanelli, L. F.; McHugh, M. A. (Methanol + ethene): phase behavior and modeling with the SAFT equation of state 625

Castells, R. C.; Menduiña, C.; Pando, C.; Renuncio, J. A. R. The excess enthalpies of (dinitrogen oxide + toluene) at the temperature 313.15 K and at pressures from 7.60 MPa to 15.00 MPa 641

Fonseca, I. M. A.; Lobo, L. Q. (Vapour + liquid) equilibrium of (methyl fluoride + dinitrogen oxide) at the temperature 182.33 K 647

Marceca, E.; Fernández-Prini, R. Solubility of solids in near-critical fluids. IV. CHI_3 and I_2 in trifluoromethane 651

McElroy, P. J. Excess second virial coefficients for (carbon monoxide + butane) and for (carbon monoxide + 2-methylpropane) 663

NOTE

Fonseca, I. M. A.; Lobo, L. Q. The triple point of methyl fluoride 671

Number 7, July 1994

Bao, J.-B.; Hang, L.-L.; Han, S.-J. Infinite-dilution activity coefficients of (propanone + an *n*-alkane) by gas stripping 673

Letcher, T. M.; Domańska, U.; Govender, P. The excess enthalpies of (di-butylamine + an ether) at the temperature 298.15 K 681

Nagata, I. Excess enthalpies for (propan-2-ol + propanone) and (propan-2-ol + propanone + benzene) at the temperature 298.15 K 691

Letcher, T. M.; Nevines, J. A. Excess volumes of (a chlorinated benzene + an alkanol) at the temperature 298.15 K 697

Artal, M.; Muñoz Embid, J.; Velasco, I. Excess enthalpies and excess volumes of (iodomethane + 1-fluoropentane or 1-chloropropane or 1-chloropentane or bromoethane or 1-bromobutane or iodoethane or 1-iodobutane) at the temperature 298.15 K 703

Nagata, I. Isothermal (vapour + liquid) equilibria of (propan-1-ol + 2-methylpropan-1-ol + acetonitrile + benzene) 709

Gospodinov, G. G. Heat capacities and thermodynamic functions of the tellurates(IV) of aluminium, gallium, indium, and thallium 713

Ott, J. B.; Sipowska, J. T.; Marchant, B. G. Excess enthalpies for (pentane + methanol) at the temperatures (283.15, 298.15, and 363.15) K and the pressures (0.4, 5, and 15) MPa 717

Tomaszkiewicz, I.; O'Hare, P. A. G. Thermochemistry of (germanium + sulfur). II. Standard molar enthalpy of formation $\Delta_f H^\circ_m$ of germanium(II) sulfide GeS at the temperature $T = 298.15$ K determined by fluorine-bomb combustion calorimetry. 727

Smruti Dash; Ziley Singh; Prasad, R.; Sood, D. D. The standard molar Gibbs free energy of formation of $Ba_3Zr_2O_7$(s). 737

Smruti Dash; Ziley Singh; Prasad, R.; Sood, D. D. The determination of the standard molar Gibbs free energy of formation of Ba_2ZrO_4(s) 745

Trusler, J. P. M. The speed of sound in $(0.8CH_4 + 0.2C_2H_6)$(g) at temperatures between 200 K and 375 K and amount-of-substance densities up to 5 mol·dm^{-3} . . 751

Joyce, D. B.; Blencoe, J. G. Excess molar Gibbs free energies for $\{xH_2O + (1-x)CO_2\}$ at temperatures from 673 K to 973 K at the pressure 50 MPa 765

Nagata, I. Excess enthalpies for (1,1-dimethylethyl methyl ether + benzene) and (methanol + 1,1-dimethylethyl methyl ether + benzene) at the temperature 298.15 K 779

Number 8, August 1994

Grolier, J.-P. E. Maria Inmaculada Paz Andrade and her scientific career . . . 785

Yang, M.-Y.; Pilcher, G.; Macnab, J. I. Standard enthalpies of formation in the crystalline state of aminomethanesulfonic acid, 2-aminoethanesulfonic acid, and the three aminobenzenesulfonic acids 787

Barbés, B.; García, I.; González, J. A.; Cobos, J. C.; Casanova, C. Excess properties of (an *n*-alkoxyethanol + an organic solvent). VI. $V^E_m\{xCH_3(CH_2)_{v-1}O(CH_2)_2O(CH_2)_2OH + (1-x)C_6H_5CH_3\}$ for $v = 1$, 2, and 4 at the temperature 298.15 K 791

Iglesias, T. P.; Legido, J. L.; Romaní, L.; Peleteiro, J.; Franjo, C. Relative permittivities, densities, and excess molar volumes of $\{xCH_3(CH_2)_4CH_3 + (1-x)CH_3(CH_2)_vOH\}$ ($v = 2$, 4, and 6) at the temperature 298.15 K 797

Pintos, M.; Amigo, A.; Bravo, R.; Calvo, E. Excess volumes of (tetrahydropyran + heptane + heptan-1-ol or octan-1-ol) at the temperature 298.15 K 803

Lorenzana, M. T.; Franjo, C.; Jiménez, E.; Baluja, M. C.; Fernández, J. Alkyl-chain-length dependence of the excess volumes for (an α,ω-dichloralkane + an alkan-2-one) 809

Jiménez, E.; Romaní, L.; Wilhelm, E.; Roux-Desgranges, G.; Grolier, J.-P. E. Excess heat capacities and excess volumes of (an *n*-alkylalkanoate + heptane or decane or toluene) 817

Cheda, J. A. R.; Fernández-Martín, F.; Burns, R. J.; Westrum, E. F., Jr. Thermodynamics of the thallium alkanoates. VII. Heat capacity and thermophysics of thallium(I) butanoate at temperatures from 6 K to 480 K 829

Ribeiro da Silva, M. A. V.; Ferrão, M. L. C. C. H.; Fang Jiye Standard enthalpies of combustion of the six dichlorophenols by rotating-bomb calorimetry . . . 839

Azevedo, E. J. S. G.; Zollweg, J. A.; Calado, J. C. G. Thermodynamics of liquid mixtures: (argon + ethane) 847

Matos, J. S.; Trenzado, J. L.; Caro, M. N.; Romano, E.; Pérez, E. Volumetric study of (an aliphatic methyl ester + heptane or nonane) at the temperature 298.15 K . 857

Berro, C.; Laïchoubi, F.; Rauzy, E. Isothermal (vapour + liquid) equilibria and excess volumes of (3-methylpentane + heptane), of (3-methylpentane + octane), and of (toluene + octane) 863

Romaní, L.; Tovar, C. A.; Carballo, E.; Peleteiro, J.; Legido, J. L. Densities and excess molar volumes of $\{xCH_3O(CH_2CH_2O)_vCH_3 + (1-x)CH_3(CH_2)_5CH_3\}$ ($v = 3, 4$) at several temperatures. 871

Rio, J. M. del; Pombo, C.; Prieto, G.; Sarmiento, F.; Mosquera, V.; Jones, M. N. *n*-Alkyltrimethylammonium bromides in a buffered medium: a thermodynamic investigation. 879

Duarte, C. M. M.; Guedes, H. J. R.; Nunes da Ponte, M. (p, V_m, T) measurements on gaseous and liquid ($0.5Xe + 0.5C_2H_6$) near the critical region 889

Fernández, J.; Williamson, A.-M.; McLure, I. A. Orthobaric surface tension of (methylcyclohexane + tetradecafluoromethylcyclohexane) in the critical region at temperatures from 320 K to 335 K 897

Number 9, September 1994

Wang Haijun; Chen Mingzhi; Zheng Guokang Excess volumes of (an aliphatic dinitrile + an aromatic hydrocarbon) at the temperature 298.15 K 913

Zielkiewicz, J. (Vapour + liquid) equilibria in (heptane + propan-2-ol or butan-1-ol or 2-methylpropan-1-ol or 2-methylpropan-2-ol or pentan-1-ol) at the temperature 313.15 K 919

Fernández-Martín, F.; Cheda, J. A. R.; Jones, L. L.; Lin, C. C.; Westrum, E. F., Jr. Thermodynamics of the thallium alkanoates. VIII. Heat capacity and derived thermophysical properties of thallium(I) propanoate and pentanoate at temperatures from 6 K to 490 K 925

Lebedev, B. V.; Kulagina, T. G.; Smirnova, N. N. Thermodynamics of butylmethacrylate and perdeuterobutylmethacrylate at temperatures from 0 to 330 K . . . 941

Zielkiewicz, J. Excess volumes of (heptane + propan-2-ol or butan-1-ol or 2-methylpropan-1-ol or 2-methylpropan-2-ol or pentan-1-ol) at the temperature 313.15 K 959

Klomfar, J.; Hrubý, J.; Šifner, O. Measurements of the (T, p, ρ) behaviour of 1,1,1,2,3,3,3-heptafluoropropane (refrigerant R227) in the liquid phase . . . 965

Masuda, N.; Nagano, Y.; Sakiyama, M. Standard molar enthalpy of formation of $(CH_3)_2SO$, dimethylsulfoxide, by combustion calorimetry 971

Aim, K. A modified ebulliometric method for high-boiling substances: vapour pressures of 2-chlorobenzonitrile and 4-chlorobenzonitrile at temperatures from 380 K to 490 K 977

Stølen, S.; Fjellvåg, H.; Grønvold, F.; Seim, H.; Westrum, E. F., Jr. Phase stability and structural properties of $Ni_{7+\delta}S_6$ and Ni_9S_8. Heat capacity and thermodynamic properties of Ni_7S_6 at temperatures from 5 K to 970 K and of Ni_9S_8 from 5 K to 673 K 987

Diky, V. V.; Kabo, G. J.; Kozyro, A. A.; Krasulin, A. P.; Sevruk, V. M. Thermodynamic properties of crystalline and liquid chlorocyclohexane and inversion of ring . 1001

Nagata, I. Excess enthalpies of (methanol + aniline + propanone) at the temperature 298.15 K 1015

Letcher, T. M.; Domańska, U.; Govender, P. The excess volumes of (dibutylamine + an ether) at the temperature 298.15 K 1019

Number 10, October 1994

Franjo, C.; Lorenzana, M. T.; Legido, J. L.; Paz Andrade, M. I.; Jiménez, E. Excess molar volumes of $[x_1CH_3CH_2CO_2(CH_2)_2CH_3 + x_2CH_3(CH_2)_5OH + (1-x_1-x_2)\{CH_3(CH_2)_4CH_3$ or $c\text{-}C_6H_{12}\}]$ at the temperature 298.15 K . . 1025

Abdulagatov, I. M.; Polikhronidi, N. G.; Batyrova, R. G. Measurements of the isochoric heat capacities C_V of carbon dioxide in the critical region 1031

Nikitin, E. D.; Pavlov, P. A.; Popov, A. P. (Vapour + liquid) critical properties and acentric factors of five low-molar-mass methylphenylsiloxanes 1047

Lagourette, B.; Daridon, J. L.; Gaubert, J. F.; Xans, P. Experimental determination of ultrasonic speeds in (methane + propane)(g) and in (methane + octane)(g) at high pressures 1051

Pal, A.; Singh, Y. P. Excess molar volumes and apparent molar volumes of $\{xH(CH_2)_2O(CH_2)_2O(CH_2)_2OH + (1-x)H_2O\}$ at the temperature 298.15 K . 1063

Christov, C. Thermodynamic study of $(b_1RbCl + b_2MeCl_2)(aq)$, where b denotes molality and Me denotes Mn, Co, Ni, or Cu, at the temperature 298.15 K, on the basis of Pitzer's model 1071

Tomaszkiewicz, I.; Susman, S.; Volin, K. J.; O'Hare, P. A. G. Fluorine-combustion calorimetric determinations of the standard molar enthalpy changes for the formation of $SiSe_2$(cr), $SiSe_{1.94}$(cr), and $SiSe_{1.94}$(vit), and for the transition: $SiSe_{1.94}$(vit) = $SiSe_{1.94}$(cr) at the temperature $T = 298.15$ K. Implications of the results for the enthalpies of dissociation D°_m(Se—SiSe) and D°_m(SiSe). Thermodynamic properties of SiSe(g) 1081

Huntelaar, M. E.; Booij, A. S.; Cordfunke, E. H. P. The standard molar enthalpies of formation of $BaZrO_3$(s) and $SrZrO_3$(s) 1095

Carson, A. S.; Laye, P. G.; Pedley, J. B.; Welsby, A. M.; Chikos, J. S.; Hosseini, S. The enthalpies of formation of iodoethane, 1,2-diiodoethane, 1,3-diiodopropane, and 1,4-diiodobutane 1103

Gospodinov, G. G. Heat capacities of copper tellurates(IV) at temperatures from 350 K to 550 K 1111

Kleut, G. J. van de; Sijpkes, A. H.; Gill, S. C. The solubilities of five cyclic dipeptides in water at the temperature 298.15 K 1115

Rodríguez-Raposo, R.; Fernández-Mérida, L.; Esteso, M. A. Activity coefficients in (electrolyte + amino acid)(aq). The dependence of the ion–zwitterion interactions on the ionic strength and on the molality of the amino acid analysed in terms of Pitzer's equations 1121

Ichihara, T.; Uematsu, M. Densities of liquid ammonia in the temperature range from 310 K to 360 K at pressures up to 200 MPa 1129

Number 11, November 1994

Nagata, I. Excess enthalpies for (ethanol + 1,1-dimethylethyl methyl ether) and (ethanol + 1,1-dimethylethyl methyl ether + benzene) at the temperature 298.15 K 1137

Majer, V.; Wood, R. H. Volumetric properties of aqueous 1–1 electrolytes near and above the critical temperature of water. III. Experimental densities and apparent molar volumes of CsBr(aq) to the temperature 725.5 K and the pressure 38.0 MPa, comparison with other 1–1 electrolytes, and extrapolations to infinite dilution for NaCl(aq) 1143

Faizullin, M. Z.; Skripov, V. P. The (T, p, x) surface of phase separation of $[(1-x)H_2O + x(CH_3)\{CH_3(CH_2)_4\}_2PO]$ in the vicinity of the lower critical solution temperatures at pressures up to 230 MPa 1167

Rodríguez, V.; Artigas, H.; Lafuente, C.; Royo, F. M.; Urieta, J. S. Excess volumes of (1,2-dichloroethane or 1,2-dibromoethane + butan-1-ol or butan-2-ol or 2-methylpropan-1-ol or 2-methylpropan-2-ol) at the temperatures 298.15 K and 313.15 K . 1173

Iglesias, M.; Orge, B.; Tojo, J. Speeds of sound, densities, refractive indices, and isentropic compressibilities of $\{x_1C_6H_6 + x_2c\text{-}C_6H_{12} + (1-x_1-x_2)C_6H_5Cl\}$ at the temperature 298.15 K 1179

Chirico, R. D.; Hossenlopp, I. A.; Gammon, B. E.; Knipmeyer, S. E.; Steele, W. V. Heat capacities of the six dimethylpyridines between the temperatures 10 K and 445 K and methyl-group rotational barriers in the solid state 1187

Chirico, R. D.; Hossenlopp, I. A.; Gammon, B. E.; Knipmeyer, S. E.; Steele, W. V. Heat capacities between the temperatures 10 K and 445 K for 2,4-dimethylpyrrole and barriers to methyl-group rotation in the solid state for 2,4- and 2,5-dimethylpyrrole 1219

Tanaka, T.; Atake, T.; Nakayama, H.; Eguchi, T.; Saito, K.; Ikemoto, I. Heat capacity and an incommensurate phase transition of crystalline bis(4-chlorophenyl)sulfone 1231

Letcher, T. M.; Domańska, U. The excess volumes of (tributylamine + an ether) at the temperature 298.15 K 1241

Number 12, December 1994

Conti, G.; Gianni, P.; Matteoli, E. Excess enthalpies and excess heat capacities of (*N*,*N*-dimethylformamide + tetrahydrofuran + cyclohexane) at the temperature 298.15 K 1249

Takai, S.; Akishige, Y.; Kawaji, H.; Atake, T.; Sawaguchi, E. Low-temperature heat capacities and Verwey transition of magnetite 1259

Zubkov, A. A. Dissolution enthalpies of 3d transition metals in liquid aluminium . 1267

Sipowska, J. T.; Lemon, L. R.; Ott, J. B.; Brown, P. R.; Marchant, B. G. Excess enthalpies and excess volumes for (butane + butan-1-ol) at the temperatures (298.15, 323.15, and 348.15) K and pressures (5, 10, and 15) MPa . . . 1275

Nagata, I. Isothermal (vapour + liquid) equilibria of (ethanol + propan-2-ol + acetonitrile + benzene) and (propan-2-ol + butan-2-ol + acetonitrile + benzene) 1287

Nagata, I. Excess enthalpies for (methanol + benzene + cyclohexane) and (methanol + benzene + heptane) at the temperature 298.15 K 1293

Jin, X. W.; Wang, Z. C. Solubility of a solute mixture: (iodine + anthracene) in benzene or toluene or each of the three xylenes at the temperature 298.15 K . . . 1297

González, E.; Ortega, J.; Postigo, M. A.; Legido, J. L. Thermodynamic properties of (a methyl ester + an *n*-alkane). VII. H_m^E and V_m^E for $\{x CH_3(CH_2)_{2u-1}CO_2CH_3 + (1-x)CH_3(CH_2)_{2v}CH_3\}$ for $u = 4$ to 7 and $v = 2$ to 7 1301

Zielkiewicz, J. Excess volumes of (*N*,*N*-dimethylformamide + ethanol + water) at the temperature 313.15 K 1317

Mier, W.; Lichtenthaler, R. N.; Roux, A. H.; Grolier, J.-P. E. Excess molar heat capacities $C_{p,m}^E$ and excess molar volumes V_m^E of $\{x_1 CH_3(CH_2)_5CH_3 + x_2 CH_3C(CH_3)_2CH_2CH(CH_3)CH_3 + x_3 CH_3C(CH_3)_2OC_2H_5 + (1-x_1-x_2-x_3)C_2H_5OH\}$(1). I. Binary and quaternary mixtures . . . 1323

Mier, W.; Lichtenthaler, R. N.; Roux, A. H.; Grolier, J.-P. E. Excess molar heat capacities $C_{p,m}^E$ and excess molar volumes V_m^E of $\{x_1 CH_3(CH_2)_5CH_3 + x_2 CH_3C(CH_3)_2CH_2CH(CH_3)CH_3 + x_3 CH_3C(CH_3)_2OC_2H_5 + (1-x_1-x_2-x_3)C_2H_5OH\}$(1). II. Ternary mixtures and prediction of quaternary values 1335

Ferloni, P.; Kenesey, C.; Westrum, E. F., Jr. Thermodynamics of alkali alkanoates. X. Heat capacities and thermodynamic properties of lithium methanoate and lithium ethanoate at temperatures from ≈5 K to 580 K 1349

INDEX TO VOLUME 26 1365